

Original Article

PKM ζ contributes in consolidation, retrieval and maintenance of amygdala dependent fear memory in rats

Bentolhoda Amirshabani^{1,2}, Mohammad Rostampoor^{1,2}, Parvin Babaei^{1,2,3*}

1. Department of Physiology, Faculty of Medicine, Guilan University of Medical Sciences, Rasht, Iran
2. Cellular & Molecular Research Center, Faculty of Medicine, Guilan University of Medical Sciences, Rasht, Iran
3. Neuroscience Research Centre, Faculty of Medicine, Guilan University of Medical Sciences, Rasht, Iran

Abstract

Introduction: Protein kinase M zeta (PKM ζ) is assumed to be actively involved in retaining long-term potentiation. The goal of this study was to investigate the role of PKM ζ in basolateral amygdala (BLA) upon acquisition, consolidation, retention and retrieval of memory using a specific inhibitor of PKM ζ .

Methods: Sixty male wistar rats underwent stereotaxic surgery and were cannulated bilaterally at the BLA nucleus. Then animals were divided into 4 groups of receiving BLA microinjection of zeta inhibitory peptide (ZIP) in different time courses: 30 min before and after training, 30 min before the testing (on the day after the learning) and 30 min after testing (but testing 10 days later). Memory was assessed using step through passive avoidance.

Results: ZIP infusion in BLA had no significant change on acquisition ($P=0.06$), however significantly impaired consolidation, retrieval and maintenance of passive avoidance memory ($P=0.012$).

Conclusion: Findings indicate that PKM ζ activity in the BLA plays an important role in retaining amygdala dependent avoidance memory interfering the process of consolidation, retrieval and maintenance of learned task.

Keywords:

Protein kinase M ζ (PKM ζ);
Learning;
Fear memory;
zeta inhibitory peptide (ZIP)

Received: 23 Feb 2017

Accepted: 29 Jun 2017

*Correspondence to:

P. Babaei

Tel: +98-9113313747

Fax: +98-13133690036

Email:

p_babaei@gums.ac.ir

Introduction

Amygdala is a component of the limbic system that plays a critical role in emotion, learning and memory, and complex behavior (McDonald and Mott, 2017). This brain region exhibits atrophy and accumulation of neurofibrillary tangles in Alzheimer disease (AD), which leads to impairment in memories of emotional events (Poulin et al., 2011). These findings show the

importance of amygdala in the formation of fear memory. On the other hand, post-traumatic stress disorder (PTSD) is assumed to result from impairments in fear, extinction and/or over-consolidation of fear memories due to amygdala dysfunction (Shin and Liberzon, 2010).

It has been known that the amygdala receives different sensory information via cortex, hippocampus and thalamus (McDonald, 1998). Among different nucleus of the amygdala, the anterior portion of the

basolateral nucleus (BLA), modulates consolidation of memories of emotionally arousing experiences through its projections to the hippocampal regions (McGaugh, 2004). These projection neurons of the BLA are glutamatergic pyramidal or GABAergic nonpyramidal neurons (McDonald, 1992).

Learning and memory were known to be mediated by different mechanisms such as strengthening of long term potentiation (LTP) (Giese, 2012), synaptogenesis (Stewart and Popov, 2012) and modulation of intrinsic excitability properties (Papoutsi et al., 2012). Studies of synaptic plasticity in the rat have shown that LTP is a candidate mechanism for memory formation in BLA (Ikegaya et al., 1996). Despite numerous theories regarding molecular pathways of memory formation and consolidation, the mechanisms underlying storage of established memory and retrieval have not been understood yet. It has been known that different kinds of synaptic plasticity require different protein kinases. In the adult mammalian brain, at least 250 protein kinases are expressed, but only a few of them participate in the mechanism underlying learning and memory (Giese and Mizuno, 2013). A number of kinases such as cAMP-dependent protein kinase A (PKA), protein kinase C (PKC) and Ca^{2+} /calmodulin-dependent protein kinases (CaMKII) have been shown to have the type of persistent activity following synaptic stimulation (Schwartz and Greenberg, 1987) and blocking these proteins affected the development of intermediate-term facilitation, rather than the maintenance of long-term potentiation (Sutton and Carew, 2000). protein kinase M zeta (PKM ζ) which is expressed only in the nervous system (Hernandez et al., 2003), is an atypical isoform of PKC due to the lack of inhibitory domain, sustains constantly active to preserve LTP and memory (Madrónal et al., 2010). A study in rats has shown a high level of PKM ζ expression in the prefrontal cortex and hippocampus after training in a delayed non match-to-sample task (Wang et al., 2014). Hara and his colleagues found higher level of PKM ζ in the hippocampal dentate gyrus of fast learner monkeys (Hara et al., 2012). Also, it has been shown that over-expression of PKM ζ in the prelimbic cortex selectively enhanced the formation of long-term fear memory (Xue et al., 2015). In contrast, aged monkeys with lower recognition function expressed a lower level of PKM ζ in the hippocampal dentate gyrus (Hara et al., 2012).

PKM ζ has been known to be translated within only 10 min in response to LTP-inducing stimuli (Hernandez et al., 2003), not only in cell bodies, but also locally in the dendritic area (Muslimov et al., 2004). Previous results have also suggested the role of the BLA in associative memories such as conditioned taste aversion (CTA) acquisition (Barot et al., 2008) which may be mediated by PKC activity (Yasoshima and Yamamoto, 1997). The role of the basolateral amygdala PKM ζ in memory formation and maintenance has not been fully understood. Administration of bilateral inhibitor of PKM ζ , ZIP, into the BLA induced retrograde amnesia for learned contextual and auditory fear responses, such as freezing and passive avoidance responses (Migues et al., 2010; Gámiz and Gallo, 2011).

Also, Tiunova and colleagues (2015) showed that PKM ζ involves in the maintenance of long term memory (LTM) for taste aversion learning in young chicks. In addition, PKM ζ has recently been observed to be down-regulated during memory extinction, reconsolidation blockade (Xue et al., 2012) and spatial familiarity in hippocampus (Moncada and Viola, 2008). Abnormal aggregations of PKM ζ was observed in and near neurofibrillary tangles in the brains of individuals with AD (Crary et al., 2006). Despite all these facts, whether or not PKM ζ , plays a role in BLA dependent fear memory remains to be clarified.

Therefore, to clarify the role of amygdala PKM ζ in the process of fear memory formation, this experiment was designed to investigate the effect of ZIP (myr-SIYRRGARRWRKL-OH) microinjections into the BLA upon acquisition, consolidation, retention and retrieval of memory.

Materials and methods

Animals

Sixty four male Wistar rats (3 months age and 180-200 g weight) were housed four per cage and fed rat chow and tap water and libitum. Room temperature was maintained at 22 ± 2 °C with a 12/12 h light/dark cycle (light on 7:00 am). All experiments were performed in accordance with the guidelines provided by the Experimental Animal Laboratory prepared by ethical committee of Guilan University of Medical Sciences, Rasht, Iran (Ethic code: IR.GUMS.REC.1394.425).

Surgery

Subjects were handled for three consecutive days in experimental room before surgery. Each rat was anesthetized with an intraperitoneal injection of ketamine hydrochloride (100 mg/kg) and xylazine (10 mg/kg), then were cannulated bilaterally with stainless steel 22-gauge cannula at the BLA using stereotaxic coordinates of (2.8 mm posterior, \pm 5.0 mm lateral, 7.2 mm ventral) relative to Bregma (Paxinos and Watson, 1986). Each rat was given a recovery period of at least one week before behavioral testing.

Drug

The PKM ζ inhibitor, ZIP (Tocris Bioscience, USA) (Serrano et al., 2005) was diluted in sterile saline to a concentration of 10 nmol/ μ l (Pastalkova et al., 2006). Animals received 1 μ l of ZIP per side of BLA according to the design of the experiment.

Experimental design

After recovery, animals were divided into 8 groups of receiving either BLA microinjection of ZIP or saline in different time courses as follows: 1) Animals received microinjection of ZIP into the BLA, 30 min before the training in step through passive avoidance task, and then were tested 24 h later in the same task. Control group received the same volume of saline into the BLA, 30 min before training and were tested 24 h later. 2) Animals received microinjection of ZIP into the BLA, 30 min after the training, time for consolidation of memory, and then were tested 24 h later. Control group received the same volume of saline into the BLA, 30 min after the training and were tested 24 h later. 3) Animals were trained in step through passive avoidance task and received either ZIP or saline microinjection 30 min before testing, on the day after training. 4) Animals were trained, ZIP or saline were injected 30 min after testing, then animals were tested 10 days later to assess the maintenance of memory (Parsons and Davis, 2011b).

Training

Memory traces are formed over time and includes number of phases including acquisition, or learning, consolidation (the labile phase during which the memory trace will be physically stored); Recall, or memory retrieval, which takes place during the re-exposition to the learning context, and finally storage

or maintenance of established memory (Quillfeldt, 2016). Acquisition in our experiment took place in one trial, when the animals were exposed to a context with delivery of electric foot shock. Here we injected ZIP 30 min before the learning in order to study the effect of PKM ζ inhibitor on acquisition processes.

In the second part of the experiment thirty minutes after learning, we injected ZIP to interfere consolidation phase; when the memory trace undertakes a labile phase and the animal is susceptible to disturbances (Dudai, 2000). Recall test was done after a 24h interval and animals received ZIP before exposing to the same context in which they acquired the task. Finally maintenance of LTM was assessed after 10 days in our experiment. It has been know that LTM can be assessed after minimum period of 6 hours, preferentially 24h interval, but larger time spans from several days, to months are possible according to the memory intensity, forgetting, extinction and reconsolidation (Quillfeldt, 2016).

Step through passive avoidance apparatus consisted of two-compartment dark/light shuttle box separating by a guillotine door. Floor of the dark compartment consisted of a stainless steel shock grid floor. Before initial of the experiment, animals were handled for three days in order to familiarize to the context and experimenter as well. Our protocol was single trial, placing rat individually in the light room for 20 seconds in order to explore the context. Then the door was opened and the rat was allowed to move freely into the dark room. Upon entry into the dark room, the door was closed and the rat was given 1mA electrical shock for 5 seconds and immediately was returned to its home cage. After 24 hours the rat was placed in the light room for testing the memory retrieval without delivery of any shock for 180 seconds. Latency to enter to the dark chamber and total time spent in the dark room were chosen as indicators of memory.

Statistics

Normality of variables was evaluated by Kolmogorov-Smirnov (KS) test in SPSS ver 22 and non-parametric mann-whitney test was used to compare means of ZIP and saline receiving groups based on the results of KS. *P*-value less than 0.05 was considered to be statistically significant and results are expressed as the means \pm SEM.

Results

Experiment 1: Effect of BLA injection of ZIP on acquisition of memory

In the first experiment, ZIP or saline was infused into the BLA, 30 minutes before training and the rats were tested 24 h later. As Figure 1 illustrates, animals receiving ZIP before training session showed no significant change in step through latency after 24 h, compared to saline receiving group (Fig. 1A), (166.25 ± 6.1 vs. 178.75 ± 1.2 sec $P=0.06$), or total time spent in the dark room (TTS), (8 ± 2.7 vs. 1.8 ± 1.3 , $P = 0.1$, Fig. 1B).

Experiment 2: Effect of BLA injection of ZIP on consolidation of memory

ZIP or saline were infused into the BLA, 30 minutes after training and the rats were tested 24 h later. Mann-Whitney test showed less step through latency (24.1 ± 7.5 vs. 178 ± 1.25 , $P = 0.012$, Fig. 2A), but more total time spent in the dark room (97.87 ± 18.1 , vs. 3 ± 1.51 sec, $P = 0.012$, Fig. 2B), in the animals receiving ZIP.

Experiment 3: Effect of BLA injection of ZIP on retrieval of memory




Fig.1. The effect of ZIP administration 30 min before training on acquisition of memory in step through passive avoidance learning. A) First latency to enter to the dark room, data is expressed as mean±SEM, $P=0.1$ vs. control group. B) Columns show mean±SEM of total time spent on dark room of retention. No significant difference was found between ZIP and saline groups ($P=0.06$, $n=8$ per each group).




Fig.2. The effect of ZIP administration 30 min after training on consolidation of memory in step through passive avoidance learning. A) First latency to enter to the dark room, data is expressed as mean±SEM (n=8) $P=0.012$ vs. control group. B) Columns show mean±SEM of total time spent on dark room of retention. $P=0.012$ indicates significant difference vs. control group. Data are expressed as mean±SD (** $P=0.012$, n=8 per each group).

Animals of the third group were trained in the first day and received ZIP, 30 minutes before testing on the day later, when molecular memory consolidation was completed. Mann-Whitney test analysis showed less step through latency (9.6 ± 3.1 vs. 176.25 ± 1.62 sec, $P = 0.012$, Fig. 3A) but more total time spent in the dark room (97.37 ± 20.4 vs. 6.6 ± 2.7 , $P = 0.012$, Fig. 3B) in the animals receiving ZIP 30 minutes before testing.

Experiment 4: Effect of BLA injection of ZIP on maintenance of memory

The PKMζ inhibitor, ZIP was injected 30 min after

retrieval test, 24 h after training (when molecular memory consolidation was completed and successfully recalled), but animals were tested 10 days later. Mann-Whitney test analysis showed less step through latency (9.6 ± 3.2 vs. 176.25 ± 1.62 , $P = 0.012$, Fig. 4A) but more TTS (74.5 ± 15.9 vs. 4 ± 1.5 sec, $P = 0.012$, Fig. 4B) in the animals receiving ZIP compared with saline receiving ones.

Discussion

The results of the present study in line with study carried out by (Shema et al., 2007; Crespo et al.,




Fig.3. The effects of ZIP infusion 24h after training on memory retrieval in step through passive avoidance learning. A) Columns show mean \pm SEM of step-down first latency of retention test. Mann-Whitney test analysis showed less step through latency, $P=0.012$. B) Total time spent in the dark room showed significant difference ($P=0.012$) in the animals receiving ZIP 30 minutes before testing compared with saline. Data are expressed as mean \pm SD (** $P=0.012$, $n=8$ per each group).

2012) on cortex, showed that injection of ZIP into the BLA, 30 minutes before training had no significant effect on the acquisition of CTA. As it has been known, acquisition of learning involves NMDA receptor activation and influx of calcium (Morris et al., 1986) to induce early phase of long term memory (E-LTP) (Bliss and Collingridge, 1993). E-LTP lasts 1–3 hours and does not require protein synthesis (Stanton et al., 2006), however memory acquisition or learned associations are transferred from a labile short-term memory state to a stable form, through memory consolidation (McGaugh, 2000). Blocking BLA PKM ζ ,

30 minutes after the training in the present study, significantly inhibited the consolidation of passive avoidance learning similar to the previous study on CTA in insular cortex of rats (Shema et al., 2007) and also in conditioning response (Serrano et al., 2008). To explain the mechanisms underlying this finding, it is important to notice that late phase of long term potentiation (L-LTP) is induced by a cascade of cellular signaling events following increase in intracellular cAMP concentration (Edelman et al., 1987), activation of PKA and transcription factors such as cAMP response element binding protein




Fig.4. The effects of ZIP infusion 24h after training on memory maintenance in step through passive avoidance learning. A) Columns show mean \pm SEM of step-down first latency of retention test which was performed 10 days after training. Mann-Whitney test showed significant difference vs. control group, $P=0.012$. B) Columns show mean \pm SEM of total time spent dark room of retention test. Mann-Whitney test showed less TTS, $P=0.012$. Data are expressed as means \pm SD (** $P=0.012$, $n=8$ per each group).

(CREB) as well (Abel et al., 1997). Finally, CREB-dependent transcription leads to the expression of a variety of downstream effector molecules that help to stabilize synaptic changes and memory consolidation (Alberini, 2005). Also L-LTP appears to be mediated by translation of PKM ζ which stabilizes AMPA receptors containing GluA2 (Sacktor, 2011). In rodents, LTP maintenance has been shown to be reversed applying PKM ζ inhibitor even one day after LTP induction (Serrano et al., 2005).

After consolidation, memory storage requires

alterations in gene transcription and protein synthesis (Reis et al., 2013). Our data in the experiment 3 showed that blocking the PKM ζ activity, 30 minutes before the memory retrieval successfully attenuated this process similar to the previous report (Serrano et al., 2008). Retrieval is generally thought to be involved in the reactivation or reconstruction of components of the neural ensembles activated during the initial learning. Indeed, proteins associated with consolidation have been implicated in memory retrieval (Murchison et al., 2004) and that AMPA

receptors activation in the amygdala during memory retrieval is necessary for expression of the conditioned response (Mamou et al., 2006).

In addition, injection of ZIP into the BLA after successful retrieval, erased avoidance memory tested on 10 days later. Similar to these findings have been reported by Tiunova et al. (2015) and Shema et al. (2007) on CTA and also Pastalkova et al on storage of spatial information. Also our findings are in agreement with data obtained from amygdala dependent fear conditioning (Migues et al., 2010; Kwapis et al., 2012) and avoidance learning (Gamiz and Gallo, 2011) indicate PKM ζ role in maintenance of emotional memory. Despite the existing literatures pointing to the role of PKM ζ in the process of memory formation and maintenance, there are some controversial data showing that some forms of memory could not be erased by inhibiting PKM ζ . These include short-term memories mediated by the hippocampus (Pastalkova et al., 2006), neocortex (Shema et al., 2009), spatial reference memory in a water maze task (Serrano et al., 2008) and olfactory fear memory (Parsons and Davis, 2011a).

Therefore, PKM ζ may be involved in the process of consolidation (Crespo et al., 2012), reestablishment (Zhang et al., 2016) and maintenance (Kwapis et al., 2012) of long term memory. To explain the mechanisms by which PKM ζ interfere these process we can address the issue to the fact that PKM ζ is located in the nucleus of neural cells as well as postsynaptic sites (Hernández et al., 2013) and subsequently leads to AMPA trafficking and Ca²⁺ influx. Then Ca²⁺ influx trigger an early and transient phase of CREB phosphorylation (Wu et al., 2001).

In addition, PKM ζ prevents internalization of GluR2-dependent AMPA receptors (Balaban et al., 2015) through interaction with proteins such as “kidney and brain expressed protein” (Yoshihama et al., 2009) and PKC1 (PICK1) (Yao et al., 2008). PKM ζ also helps to sustain histone acetylation levels during memory maintenance and using PKM ζ inhibitor erases long-term memory in part by reducing histone acetylation level (Ko et al., 2016). Future investigations with a design of reconsolidation and reminder in different times window of memory formation are needed to clarify the exact role of PKM ζ in learning and memory.

Finally, in clinical importance, it is suggested that inhibiting PKM ζ could provide a strategy for treatment

of drugs addictions, substance abuse, fear and PTSD and enhancing PKM ζ level might be a candidate to prevent cognition deficit in Alzheimer diseases.

Conclusion

The results indicate the contribution of PKM ζ in consolidation, retrieval and maintenance, but not acquisition of amygdala – dependent long-term fear memory in rat.

Acknowledgments

This research was supported by a grant from Research department of Guilan University Medical Sciences, Rasht, Iran.

Conflict of interest

Authors declare no conflict of interest

References

- Abel T, Nguyen PV, Barad M, Deuel TA, Kandel ER, Bourchouladze R. Genetic demonstration of a role for pka in the late phase of ltp and in hippocampus-based long-term memory. *Cell* 1997; 88: 615-626.
- Alberini CM. Mechanisms of memory stabilization: Are consolidation and reconsolidation similar or distinct processes? *Trends Neurosci* 2005; 28: 51-56.
- Balaban PM, Roshchin M, Timoshenko AK, Zuzina AB, Lemak M, Ierusalimsky VN, et al. Homolog of protein kinase m ζ maintains context aversive memory and underlying long-term facilitation in terrestrial snail helix. *Front Cell Neurosci* 2015; 9: 222.
- Barot SK, Kyono Y, Clark EW, Bernstein IL. Visualizing stimulus convergence in amygdala neurons during associative learning. *Proc Natl Acad Sci U S A* 2008; 105: 20959-20963.
- Bliss TV, Collingridge GL. A synaptic model of memory: Long-term potentiation in the hippocampus. *Nature* 1993; 361: 31-9.
- Crary JF, Shao CY, Mirra SS, Hernandez AI, Sacktor TC. Atypical protein kinase c in neurodegenerative disease i: Pkm ζ aggregates with limbic neurofibrillary tangles and ampa receptors in alzheimer disease. *J Neuropathol Exp Neurol* 2006; 65: 319-326.
- Crespo JA, Stöckl P, Ueberall F, Jenny M, Saria A, Zernig G. Activation of pkc ζ and pkm ζ in the nucleus accumbens core is necessary for the retrieval, consolidation and reconsolidation of drug memory. *PLoS one* 2012; 7: e30502.
- Edelman AM, Blumenthal DK, Krebs EG. Protein serine/threonine kinases. *Annu Rev Biochem* 1987; 56:

- 567-613.
- Gámiz F, Gallo M. Intra-amygdala zip injections impair the memory of learned active avoidance responses and attenuate conditioned taste-aversion acquisition in rats. *Learn Mem* 2011; 18: 529-533.
- Giese KP. Long-term potentiation and memory. *Memory mechanisms of health of disease* (ed. Giese KP) 2012: 1-17.
- Giese KP, Mizuno K. The roles of protein kinases in learning and memory. *Learn Mem* 2013; 20: 540-552.
- Hara Y, Punsoni M, Yuk F, Park CS, Janssen WG, Rapp PR, et al. Synaptic distributions of glua2 and pkm ζ in the monkey dentate gyrus and their relationships with aging and memory. *J Neurosci* 2012; 32: 7336-7344.
- Hernandez AI, Blace N, Crary JF, Serrano PA, Leitges M, Libien JM, et al. Protein kinase m ζ synthesis from a brain mrna encoding an independent protein kinase c ζ catalytic domain implications for the molecular mechanism of memory. *J Biol Chem* 2003; 278: 40305-40316.
- Hernández AI, Oxberry WC, Crary JF, Mirra SS, Sacktor TC. Cellular and subcellular localization of pkm ζ . *Philos Trans R Soc Lond B Biol Sci* 2013; 369: 20130140.
- Ikegaya Y, Saito H, Abe K. The basomedial and basolateral amygdaloid nuclei contribute to the induction of long-term potentiation in the dentate gyrus in vivo. *Eur J Neurosci* 1996; 8: 1833-1839.
- Ko HG, Kim Ji, Sim SE, Kim T, Yoo J, Choi SL, et al. The role of nuclear pkm ζ in memory maintenance. *Neurobiol Learn Mem* 2016; 135: 50-56.
- Kwapis JL, Jarome TJ, Gilmartin MR, Helmstetter FJ. Intra-amygdala infusion of the protein kinase m ζ inhibitor zip disrupts foreground context fear memory. *Neurobiol Learn Mem* 2012; 98: 148-153.
- Madroñal N, Gruart A, Sacktor TC, Delgado-García JM. Pkm ζ inhibition reverses learning-induced increases in hippocampal synaptic strength and memory during trace eyeblink conditioning. *PLoS one* 2010; 5: e10400.
- Mamou CB, Gamache K, Nader K. Nmda receptors are critical for unleashing consolidated auditory fear memories. *Nat Neurosci* 2006; 9: 1237-1239.
- McDonald A. Cell types and intrinsic connections of the amygdala. *The amygdala: neurobiological aspects of emotion, memory, and mental dysfunction* 1992: 67-96.
- McDonald AJ. Cortical pathways to the mammalian amygdala. *Progr Neurobiol* 1998; 55: 257-332.
- McDonald AJ, Mott DD. Functional neuroanatomy of amygdalohippocampal interconnections and their role in learning and memory. *J Neurosci Res* 2017; 95: 797-820.
- McGaugh JL. Memory--a century of consolidation. *Science* 2000; 287: 248-251.
- McGaugh JL. The amygdala modulates the consolidation of memories of emotionally arousing experiences. *Annu Rev Neurosci* 2004; 27: 1-28.
- Migues PV, Hardt O, Wu DC, Gamache K, Sacktor TC, Wang YT, et al. Pkm [zeta] maintains memories by regulating glur2-dependent ampa receptor trafficking. *Nat Neurosci* 2010; 13: 630-634.
- Moncada D, Viola H. Pkm ζ inactivation induces spatial familiarity. *Learn Mem* 2008; 15: 810-814.
- Morris R, Anderson E, Lynch Ga, Baudry M. Selective impairment of learning and blockade of long-term potentiation by an n-methyl-d-aspartate receptor antagonist, AP5. *Nature* 1986; 319: 774-776.
- Murchison CF, Zhang XY, Zhang WP, Ouyang M, Lee A, Thomas SA. A distinct role for norepinephrine in memory retrieval. *Cell* 2004; 117: 131-143.
- Muslimov IA, Nimrich V, Hernandez AI, Tcherepanov A, Sacktor TC, Tiedge H. Dendritic transport and localization of protein kinase m ζ mrna implications for molecular memory consolidation. *J Biol Chem* 2004; 279:52613-22.
- Papoutsis A, Sidiropoulou K, Poirazi P. Memory beyond synaptic plasticity: The role of intrinsic neuronal excitability. *Memory mechanisms in health and disease* (ed. Giese KP) 2012: 53-80.
- Parsons RG, Davis M. Corrigendum: Temporary disruption of fear-potentiated startle following pkm [zeta] inhibition in the amygdala. *Nat Neurosci* 2011a; 14: 295-6.
- Parsons RG, Davis M. Temporary disruption of fear-potentiated startle following pkm [zeta] inhibition in the amygdala. *Nat Neurosci* 2011b; 14: 295-296.
- Pastalkova E, Serrano P, Pinkhasova D, Wallace E, Fenton AA, Sacktor TC. Storage of spatial information by the maintenance mechanism of ltp. *Science* 2006; 313: 1141-1144.
- Paxinos G, Watson C. *The rat brain in stereotaxic coordinates* 2nd ed. Academic: New York, NY, USA 1986.
- Poulin SP, Dautoff R, Morris JC, Barrett LF, Dickerson BC, Initiative AsDN. Amygdala atrophy is prominent in early alzheimer's disease and relates to symptom severity. *Psychiatry Res* 2011; 194: 7-13.
- Quillfeldt JA. Behavioral methods to study learning and memory in rats. *Rodent model as tools in ethical biomedical research*: Springer, 2016: 271-311.
- Reis DS, Jarome TJ, Helmstetter FJ. Memory formation for trace fear conditioning requires ubiquitin-proteasome mediated protein degradation in the prefrontal cortex. *Front Behav Neurosci* 2013; 7.
- Sacktor TC. How does pkm ζ maintain long-term memory? *Nat Rev Neurosci* 2011; 12: 9-15.
- Schwartz JH, Greenberg SM. Molecular mechanisms for memory: Second-messenger induced modifications of protein kinases in nerve cells. *Ann rev Neurosci* 1987; 10: 459-476.
- Serrano P, Friedman EL, Kenney J, Taubenfeld SM,

- Zimmerman JM, Hanna J, et al. Pkm ζ maintains spatial, instrumental, and classically conditioned long-term memories. *PLoS Biol* 2008; 6: 2698-706.
- Serrano P, Yao Y, Sacktor TC. Persistent phosphorylation by protein kinase m ζ maintains late-phase long-term potentiation. *J Neurosci* 2005; 25: 1979-1984.
- Shema R, Hazvi S, Sacktor TC, Dudai Y. Boundary conditions for the maintenance of memory by pkm ζ in neocortex. *Learn Mem* 2009; 16: 122-128.
- Shema R, Sacktor TC, Dudai Y. Rapid erasure of long-term memory associations in the cortex by an inhibitor of pkm ζ . *Science* 2007; 317: 951-953.
- Shin LM, Liberzon I. The neurocircuitry of fear, stress, and anxiety disorders. *Neuropsychopharmacology* 2010; 35: 169-191.
- Stanton PK, Bramham C, Scharfman HE. *Synaptic plasticity and transsynaptic signaling*: Springer Science & Business Media, 2006.
- Stewart MG, Popov VI. Structural synaptic and dendritic spine plasticity in the hippocampus. *Memory mechanisms in health and disease* (ed. Giese KP) 2012: 19-52.
- Sutton MA, Carew TJ. Parallel molecular pathways mediate expression of distinct forms of intermediate-term facilitation at tail sensory-motor synapses in aplysia. *Neuron* 2000; 26: 219-231.
- Tiunova A, Bezryadov D, Anokhin K. Involvement of protein kinase m ζ in the maintenance of long-term memory for taste aversion learning in young chicks. *Bull Exp Biol Med* 2015; 158: 592-594.
- Wang DC, Liu PC, Hung HS, Chen TJ. Both pkm ζ and kibra are closely related to reference memory but not working memory in a t-maze task in rats. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 2014; 200: 77-82.
- Wu GY, Deisseroth K, Tsien RW. Activity-dependent creb phosphorylation: convergence of a fast, sensitive calmodulin kinase pathway and a slow, less sensitive mitogen-activated protein kinase pathway. *Proc Natl Acad Sci U S A* 2001; 98: 2808-13.
- Xue YX, Luo YX, Wu P, Shi HS, Xue LF, Chen C, et al. A memory retrieval-extinction procedure to prevent drug craving and relapse. *Science* 2012; 336: 241-245.
- Xue YX, Zhu ZZ, Han HB, Liu JF, Meng SQ, Chen C, et al. Overexpression of protein kinase m ζ in the prelimbic cortex enhances the formation of long-term fear memory. *Neuropsychopharmacology* 2015; 40: 2146-2156.
- Yao Y, Kelly MT, Sajikumar S, Serrano P, Tian D, Bergold PJ, et al. Pkm ζ maintains late long-term potentiation by n-ethylmaleimide-sensitive factor/gluR2-dependent trafficking of postsynaptic ampa receptors. *J Neurosci* 2008; 28: 7820-7827.
- Yasoshima Y, Yamamoto T. Rat gustatory memory requires protein kinase c activity in the amygdala and cortical gustatory area. *Neuroreport* 1997; 8: 1363-1367.
- Yoshihama Y, Hirai T, Ohtsuka T, Chida K. Kibra co-localizes with protein kinase m ζ (pkm ζ) in the mouse hippocampus. *Biosci Biotechnol Biochem* 2009; 73: 147-151.
- Zhang Y, Zong W, Zhang L, Ma Y, Wang J. Protein kinase m ζ and the maintenance of long-term memory. *Neurochem Int* 2016; 99: 215-220.